

Effective chiral theory of mesons, coefficients of ChPT, axial-vector symmetry breaking

Bing An Li

Department of Physics and Astronomy, University of Kentucky

Lexington, KY 40506, USA

Abstract

A phenomenological successful effective chiral theory of pseudoscalar, vector, and axial-vector mesons is introduced. Based on this theory all the 10 coefficients of ChPT are predicted. It has been found that a new symmetry breaking-axial-vector symmetry breaking is responsible for the mass difference between ρ and a_1 mesons. It is shown that in an EW theory without Higgs m_W and m_Z are dynamically generated by the combination of the fermion masses and the axial-vector symmetry breaking: $m_W = \frac{1}{\sqrt{2}}gm_t$, $m_Z = \rho m_W / \cos\theta_W$ with $\rho \approx 1$, and $G_F = 1/(2\sqrt{2}m_t^2)$. Two gauge fixing terms of W and Z fields are dynamically generated too.

1 Effective chiral theory of mesons

The chiral perturbation theory is rigorous and phenomenologically successful in describing the physics of the pseudoscalar mesons at low energies($E < m_\rho$). Models attempt to deal with the two main frustrations that the ChPT is limited to pseudoscalar mesons at low energy and contains many coupling constants which must be measured.

I have proposed an effective chiral theory of pseudoscalar, vector, and axial-vector mesons[1]. It provides a unified study of pseudoscalar, vector, and axial-vector meson physics at low energies.

The ansatz made in this theory is that the meson fields are simulated by quark operators. For example,

$$\rho_\mu^i = -\frac{1}{g_\rho m_\rho^2} \bar{\psi} \tau_i \gamma_\mu \psi. \quad (1)$$

The ansatz can be tested. Applying PCAC, current algebra, and this expression to the decay $\rho \rightarrow \pi\pi$, under the soft pion approximation it is derived

$$\frac{1}{2} f_{\rho\pi\pi} g_\rho = 1 \quad (2)$$

which is just the result of the VMD.

As a matter of fact, using fermion operator to simulate a meson field has a long history:

1. More than six decades ago, Jordan et al observed

$$\frac{1}{\sqrt{\pi}}\partial_\mu\phi = \bar{\psi}\gamma_5\gamma_\mu\psi \quad (3)$$

in 1 + 1 field theory.

2. A similar relation is proved in the bosonization of 1+1 field theory.
3. Use of quark operators to simulate meson fields has been already exploited in the Nambu-Jona-Lasinio(NJL) model, a model of four quark interactions.
4. Quark operators have been taken as interpolating fields in lattice gauge calculations.

The simulations of the meson fields by quark operators are realized by a Lagrangian which is constructed by chiral symmetry

$$\begin{aligned} \mathcal{L} = & \bar{\psi}(x)(i\gamma \cdot \partial + \gamma \cdot v + \gamma \cdot a\gamma_5 - mu(x))\psi(x) \\ & -\psi(\bar{x})M\psi(x) + \frac{1}{2}m_0^2(\rho_i^\mu\rho_{\mu i} + K^{*\mu}K_\mu^* + \omega^\mu\omega_\mu + \phi^\mu\phi_\mu \\ & + a_i^\mu a_{\mu i} + K_1^\mu K_{1\mu} + f^\mu f_\mu + f_{1s}^\mu f_{1s\mu}) \end{aligned} \quad (4)$$

$u = \exp\{i\gamma_5(\tau_i\pi_i + \lambda_a K_a + \lambda_8\eta_8 + \eta_0)\}$. To avoid double counting, there are no kinetic terms of mesons, which are generated dynamically. Using the least action principle, the relationships between meson fields and quark operators are found from the Lagrangian. Taking the case

of two flavors as an example, from the least action principle we obtain

$$\begin{aligned}
\frac{\Pi_i}{\sigma} &= i(\bar{\psi}\tau_i\gamma_5\psi + ix\bar{\psi}\tau_i\psi)/(\bar{\psi}\psi + ix\bar{\psi}\gamma_5\psi), \\
x &= (i\bar{\psi}\gamma_5\psi - \frac{\Pi_i}{\sigma}\bar{\psi}\tau_i\psi)/(\bar{\psi}\psi + i\frac{\Pi_i}{\sigma}\bar{\psi}\tau_i\gamma_5\psi), \\
\rho_\mu^i &= -\frac{1}{m_0^2}\bar{\psi}\tau_i\gamma_\mu\psi, \quad a_\mu^i = -\frac{1}{m_0^2}\bar{\psi}\tau_i\gamma_\mu\gamma_5\psi, \\
\omega_\mu &= -\frac{1}{m_0^2}\bar{\psi}\gamma_\mu\psi, \quad f\mu = -\frac{1}{m_0^2}\bar{\psi}\gamma_\mu\gamma_5\psi,
\end{aligned} \tag{5}$$

where $\sigma + i\gamma_5\tau \cdot \Pi = ue^{-i\eta\gamma_5}$, $\sigma = \sqrt{1 - \Pi^2}$, and $x = \tan\eta$. The pseudoscalar fields have very complicated quark structures. Substituting these expressions into the Lagrangian of two flavors, a Lagrangian of quarks is obtained. It is no longer a theory of four quarks. This model is different from NJL model.

Using path integral to integrate out the quark fields, the effective Lagrangian of mesons is derived.

$$\mathcal{L}_E = \ln \det \mathcal{D},$$

$$\mathcal{L}_{re} = \frac{1}{2} \ln \det (\mathcal{D}^\dagger \mathcal{D}),$$

$$\mathcal{L}_{im} = \frac{1}{2} \ln \det (\mathcal{D}/\mathcal{D}^\dagger).$$

The question is whether the masses, decay widths, and interactions of mesons can be described by the quark operators correctly. Taking masses as an example,

1. Pseudoscalar mesons

$$\begin{aligned}
m_{\pi^\pm}^2 &= \frac{4}{f_{\pi^\pm}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_d) - \frac{F^2}{4} (m_u + m_d)^2 \right\}, \\
m_{\pi^0}^2 &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_d) - \frac{F^2}{2} (m_u^2 + m_d^2) \right\}, \\
m_{K^+}^2 &= \frac{4}{f_{K^+}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_s) - \frac{F^2}{4} (m_u + m_s)^2 \right\}, \\
m_{K^0}^2 &= \frac{4}{f_{K^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_d + m_s) - \frac{F^2}{4} (m_d + m_s)^2 \right\}, \\
m_{\eta_8}^2 &= \frac{4}{f_{\eta_8}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle \frac{1}{3} (m_u + m_d + 4m_s) \right. \\
&\quad \left. - \frac{F^2}{6} (m_u^2 + m_d^2 + 4m_s^2) \right\}, \tag{6}
\end{aligned}$$

The formulas at the first order in quark masses are Gell-Mann, Oakes, and Renner chiral perturbation theory.

2. Vector mesons

$$m_\rho^2 = m_\omega^2 = 6m^2[2].$$

3. Axial-vector mesons

$$\begin{aligned}
\left(1 - \frac{1}{2\pi^2 g^2}\right) m_a^2 &= 6m^2 + m_\rho^2, \\
\left(1 - \frac{1}{2\pi^2 g^2}\right) m_f^2 &= m_\rho^2 + m_\omega^2,
\end{aligned}$$

The masses of vector originate in dynamical chiral symmetry breaking. This is the reason why they are much heavier than pseudoscalars. The masses of the axial-vector mesons are

resulted by a new symmetry breaking-axial-vector symmetry breaking.

The widths are

$$\Gamma_\rho = 142 MeV (Exp. 150 MeV),$$

$$\Gamma_a = 386 MeV (exp. \sim 400 MeV),$$

$$\Gamma_\omega = 7.7 MeV (exp. 7.49 MeV),$$

$$\Gamma_{f \rightarrow \rho \pi \pi} = 6.01 MeV (exp. 6.96(1 \pm 0.33) MeV)$$

This theory has following features:

1. The theory is chiral symmetric in the limit of $m_q \rightarrow 0$. The theory has dynamically chiral symmetry breaking(m),
2. VMD is a natural result

$$\frac{e}{f_v} \left\{ -\frac{1}{2} F^{\mu\nu} (\partial_\mu \rho_\nu - \partial_\nu \rho_\mu) + A^\mu j_\mu \right\}.$$

3. Axial-vector currents are bosonized

$$-\frac{g_W}{4f_a} \frac{1}{f_a} \left\{ -\frac{1}{2} F^{i\mu\nu} (\partial_\mu a_\nu^i - \partial_\nu a_\mu^i) + A^{i\mu} j_\mu^i \right\}$$

$$-\frac{g_W}{4} \Delta m^2 f_a A_\mu^i a^{i\mu} - \frac{g_W}{4} f_\pi A^{i\mu} \partial_\mu \pi^i,$$

Axial-vector symmetry breakin is taken part in.

4. The Wess-Zumino-Witten anomalous action is the leading term of the imaginary part of the effective Lagrangian,
5. Weinberg's first sum rule is satisfied analytically,
6. The constituent quark mass is introduced as m ,
7. Theoretical results of the masses and strong, E&M, and weak decay widths of mesons agree well with data,
8. The form factors of pion, π_{l3} , K_{l3} , $\pi \rightarrow e\gamma\nu$, and $K \rightarrow e\gamma\nu$ are obtained and agree with data. For example,

$$\langle r^2 \rangle_\pi = 0.445 fm^2,$$

$$Exp. = 0.44 \pm 0.01 fm^2,$$

$$\rho - pole = 0.39 fm^2.$$

9. The theory has been applied to τ mesonic decays[3] successfully. For example,

$$B(\tau \rightarrow \eta 3\pi\nu) = 3.4 \times 10^{-4},$$

$$Exp. = (4.1 \pm 0.7 \pm 0.7) \times 10^{-4},$$

$$ChPT = 1.2 \times 10^{-6}.$$

10. $\pi\pi$ and πK scatterings are studied. Theory agrees with data,
11. The parameters of this theory are: m (quark condensate), g (universal coupling constant), and three current quark masses,
12. Large N_C expansion is natural in this theory. All loop diagrams of mesons are at higher orders in N_C expansion. So far all calculations are done at the three level,
13. A cut-off has been determined to be 1.6GeV. All the masses of mesons are below the cut-off. The theory is self consistent,

2 Coefficients of chiral perturbation theory

Chiral perturbation theory is the low energy limit of any successful effective meson theory

$$\begin{aligned}
\mathcal{L} = & \frac{f_\pi^2}{16} \text{Tr} D_\mu U D^\mu U^\dagger + \frac{f_\pi^2}{16} \text{Tr} \chi (U + U^\dagger) \\
& + L_1 [\text{Tr} (D_\mu U D^\mu U^\dagger)]^2 + L_2 (\text{Tr} D_\mu U D_\nu U^\dagger)^2 \\
& + L_3 \text{Tr} (D_\mu U D^\mu U^\dagger)^2 + L_4 \text{Tr} (D_\mu U D^\mu U^\dagger) \text{Tr} \chi (U + U^\dagger) \\
& + L_5 \text{Tr} D_\mu U D^\mu U^\dagger (\chi U^\dagger + U \chi) + L_6 [\text{Tr} \chi (U + U^\dagger)]^2 \\
& + L_7 [\text{Tr} \chi (U - U^\dagger)]^2 + L_8 \text{Tr} (\chi U \chi U + \chi U^\dagger \chi U^\dagger) \\
& - i L_9 \text{Tr} (F_{\mu\nu}^L D^\mu U D^\nu U^\dagger + F_{\mu\nu}^R D^\mu U^\dagger D^\nu U)
\end{aligned}$$

$$+L_{10}Tr(F_{\mu\nu}^L U F^{\mu\nu R} U^\dagger). \quad (7)$$

Many models try to predict the coefficients of ChPT (see Table I)

The effective chiral theory of mesons is used to predict all the 10 coefficients.

1. The effective L of $\pi\pi$ scattering at low energy is derived. Two of the three coefficients are obtained

$$\begin{aligned} 2(L_1 + L_2) + L_3 &= \frac{1}{4} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^4 \\ L_2 &= \frac{1}{4} \frac{c^4}{g^2} + \frac{1}{4} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{4c^2}{g^2}\right) \\ &\quad + \frac{1}{8} \left(1 - \frac{2c}{g}\right) \frac{c}{g} \left\{2gc + \frac{1}{\pi^2} \left(1 - \frac{2c}{g}\right)\right\} \\ c &= \frac{f_\pi^2}{2gm_\rho^2}. \end{aligned} \quad (8)$$

2. A complete determination of $L_{1,2,3}$ is carried out from the effective L of πK scattering

$$\begin{aligned} -2L_1 + L_2 &= 0, \\ L_3 &= -\frac{3}{16} \frac{2c}{g} \left(1 - \frac{2c}{g}\right) \left\{2gc + \frac{1}{\pi^2} \left(1 - \frac{2c}{g}\right)\right\} \\ &\quad - \frac{1}{2} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{8c^2}{g^2}\right) - \frac{3}{4} \frac{c^4}{g^2}, \\ L_1 &= \frac{1}{32} \frac{2c}{g} \left(1 - \frac{2c}{g}\right) \left\{2gc + \frac{1}{\pi^2} \left(1 - \frac{2c}{g}\right)\right\} \\ &\quad + \frac{1}{8} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{4c^2}{g^2}\right) + \frac{1}{4} \frac{c^4}{g^2}. \end{aligned} \quad (9)$$

3. The coefficients L_{4-8} are determined from the quark mass expansions of m_π^2 , m_K^2 , m_η^2 , f_π , f_K , and f_η

$$L_4 = 0, \quad L_6 = 0, \quad (10)$$

$$L_5 = \frac{f_{\pi 0}^2 f}{8mB_0}, \quad (11)$$

$$L_8 = -\frac{F^2}{16B_0^2}, \quad (12)$$

$$3L_7 + L_8 = -\frac{F^2}{16B_0^2}, \quad L_7 = 0, \quad (13)$$

where

$$B_0 = \frac{4}{f_{\pi 0}^2} \left(-\frac{1}{3}\right) < \bar{\psi}\psi > . \quad (14)$$

L_5 and L_8 are written as

$$L_5 = \frac{1}{32Q} \left(1 - \frac{2c}{g}\right) \left\{ \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{1}{2\pi^2 g^2}\right) - \left(1 - \frac{2c}{g}\right) + \frac{4}{\pi^2} Q \left(1 - \frac{c}{g}\right) \right\}, \quad (15)$$

$$L_8 = -\frac{1}{1536g^2Q^2} \left(1 - \frac{2c}{g}\right)^2, \quad (16)$$

where

$$Q = -\frac{1}{108g^4} \frac{1}{m^3} < \bar{\psi}\psi > . \quad (17)$$

Q is a function of the universal coupling constant g only. By fitting $\rho \rightarrow e^+e^-$, g is determined to be 0.39. The numerical value of Q is 4.54.

4. L_9 and L_{10} are determined by $\langle r^2 \rangle_\pi$ and the amplitudes of pion radiative decay,

$$\pi^- \rightarrow e^- \gamma \nu$$

$$L_9 = \frac{f_\pi^2}{48} \langle r_\pi^2 \rangle, \quad (18)$$

$$L_9 = \frac{1}{32\pi^2} \frac{R}{F^V}, \quad (19)$$

$$L_{10} = \frac{1}{32\pi^2} \frac{F^A}{F^V} - L_9, \quad (20)$$

$$L_9 = \frac{1}{4}cg + \frac{1}{16\pi^2} \left\{ \left(1 - \frac{2c}{g}\right)^2 - 4\pi^2 c^2 \right\}, \quad (21)$$

$$L_{10} = -\frac{1}{4}cg + \frac{1}{4}c^2 - \frac{1}{32\pi^2} \left(1 - \frac{2c}{g}\right)^2. \quad (22)$$

There are two parameters: g and f_π^2/m_ρ^2 in all the 10 coefficients, which have been determined already. The numerical values of the 10 coefficients predicted by present theory are shown in Table II.

3 Axial-vector symmetry breaking and W and Z masses

Pion, ρ and a_1 are made of u and d quarks. Pions are Goldstone bosons. π is light. m_ρ is resulted in dynamical chiral symmetry breaking (quark condensate) and is much heavier than pions. It is well known that a_1 meson is the chiral partner of ρ meson. However, a_1 is much heavier (1.26 GeV) than ρ meson (0.77 GeV) is.

Why?

What kind of symmetry breaking is responsible for the mass difference between ρ and a_1 mesons?

There must be a new dynamical symmetry breaking which generates the mass difference between a_1 and ρ . It has been found out that the axial-vector coupling results a new dynamical symmetry breaking -axial-vector symmetry breaking which leads to

$$(1 - \frac{1}{2\pi^2 g^2})m_a^2 = 6m^2 + m_\rho^2.$$

Can this axial-vector symmetry breaking bring something to the EW theory?

A Lagrangian of EW interactions without Higgs is studied[4]

$$\begin{aligned}\mathcal{L} = & -\frac{1}{4}A_{\mu\nu}^i A^{i\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu} + \bar{q}\{i\gamma \cdot \partial - M\}q \\ & + \bar{q}_L\{\frac{g}{2}\tau_i\gamma \cdot A^i + g'\frac{Y}{2}\gamma \cdot B\}q_L + \bar{q}_R g'\frac{Y}{2}\gamma \cdot B q_R \\ & + \bar{l}\{i\gamma \cdot \partial - M_f\}l + \bar{l}_L\{\frac{g}{2}\tau_i\gamma \cdot A^i - \frac{g'}{2}\gamma \cdot B\}l_L \\ & - \bar{l}_R g'\gamma \cdot B l_R.\end{aligned}\tag{23}$$

There are no $m_{W,Z}$ and there are fermion mass terms which break the charged part of the $SU(2)_L \times U(1)$ symmetry. This explicit symmetry breaking makes charged bosons, W, massive. However, there are still two U(1) symmetries. Z and photon are massless. In EW interactions there are axial-vector couplings, therefore, there are axial-vector symmetry breaking which contribute mass to both W and Z bosons. Using

path integral to integrate out the fermion fields, the Lagrangian of boson fields is derived.

After multiplicative renormalization following results are obtained

1.

$$m_W^2 = \frac{1}{2}g^2\{m_t^2 + m_b^2 + m_c^2 + m_s^2 + m_u^2 + m_d^2 + m_{\nu_e}^2 + m_e^2 + m_{\nu_\mu}^2 + m_\mu^2 + m_{\nu_\tau}^2 + m_\tau^2\} \quad (24)$$

$$m_W = \frac{g}{\sqrt{2}}m_t. \quad (25)$$

Using the values $g = 0.642$ and $m_t = 180 \pm 12 \text{ GeV}$, it is found

$$m_W = 81.71(1 \pm 0.067) \text{ GeV}, \quad (26)$$

which is in excellent agreement with data $80.33 \pm 0.15 \text{ GeV}$.

2.

$$G_F = \frac{1}{2\sqrt{2}m_t^2} = 0.96 \times 10^{-5}(1 \pm 0.13)m_N^{-2}.$$

3.

$$m_Z^2 = \rho m_W^2 / \cos^2 \theta_W, \quad (27)$$

$$\rho = (1 - \frac{\alpha}{4\pi}f_4)^{-1},$$

$$f_4 = \frac{1}{3}N_G - \frac{2}{3}\sum_q f_3 + \frac{2}{3}\sum_l f_3.$$

$$f_3 = \frac{1}{2} \frac{1}{\sqrt{x}} \ln \frac{1+\sqrt{x}}{1-\sqrt{x}}.$$

$$x = \left(\frac{m_2^2}{m_1^2}\right)^2, \quad m_{1,2} = \frac{1}{2}(m_t \pm m_b).$$

$$\rho \sim 1,$$

$$m_Z^2 = m_W^2 / \cos^2 \theta_W.$$

4. The propagator of W field is derived

$$\frac{i}{q^2 - m_W^2} \{-g_{\mu\nu} + \frac{q_\mu q_\nu}{q^2}\} - \frac{i}{\xi_W q^2 - m_W^2} \frac{q_\mu q_\nu}{q^2}. \quad (28)$$

Changing the index W to Z, the propagator of Z boson field is obtained.

$$\xi_W = \frac{N_G}{(4\pi)^2} \frac{g^2}{3}, \quad \xi_Z = \frac{N_G}{(4\pi)^2} \left(1 - \frac{\alpha}{4\pi} f_4\right)^{-1} \frac{1}{3} (g^2 + g'),$$

$$N_G = 3N_C + 3.$$

1. $m_{W,Z}$ are correctly dynamically generated by axial-vector symmetry bre
2. The propagators of W and Z fields have no problems with renormalization.

4 Conclusions

1. The effective chiral theory is phenomenologically successful,
2. All the 10 coefficients of ChPT are predicted,

3. What can we learned about QCD from this theory:
 - (a) Chiral symmetry,
 - (b) Dynamical chiral symmetry breaking,
 - (c) Large N_C expansion,
 - (d) Simulations of meson fields by quark operators are working. It is worth to do theoretical study on whether the simulations are solutions of QCD at least in the limit of large N_C expansion,
4. The Lagrangian is not closed. The quark condensate, m , should be related to gluons,
5. Axial-vector symmetry breaking provides an explanation to the mass difference between ρ and a_1 ,
6. $m_{W,Z}$ are dynamically generated from the fermion masses and the axial-vector symmetry breaking.

References

- [1] B.A.Li, Phys. Rev. **D52**,5165(1995);5184(1995).

- [2] B.A.Li, Proc. of Intern. Europhys. Conf. on High Energy Phys., Brussels, Belgium, 27 Jul-2 Aug, 1995, edited by J.Lemonne *et al.*, p.225.
- [3] B.A.Li, Phys.Rev. **D55**,1425(1997).
- [4] B.A.Li, hep-ph/9711397.

Table 1: Coefficients obtained by various models

	Vectors	Quark			Nucleon loop	Linear σ model	ENJL
$(L_1 + \frac{1}{2}L_3) \times 10^{-3}$	-2.1	-.8	2.1	1.1	-.8	-.5	
$L_1 \times 10^{-3}$							0.8
$L_2 \times 10^{-3}$	2.1	1.6	1.6	1.8	.8	1.5	1.6
$L_3 \times 10^{-3}$							-4.1
$L_4 \times 10^{-3}$							0.
$L_5 \times 10^{-3}$							1.5
$L_6 \times 10^{-3}$							0.
$L_7 \times 10^{-3}$							
$L_8 \times 10^{-3}$							0.8
$L_9 \times 10^{-3}$	7.3	6.3	6.7	6.1	3.3	.9	6.7
$L_{10} \times 10^{-3}$	-5.8	-3.2	-5.8	-5.2	-1.7	-2.0	-5.5

Table 2: Coefficients from this theory

$10^3 L_1$	$10^3 L_2$	$10^3 L_3$	$10^3 L_4$	$10^3 L_5$	$10^3 L_6$	$10^3 L_7$	$10^3 L_8$	$10^3 L_9$	$10^3 L_{10}$
$.9 \pm .3$	$1.7 \pm .7$	-4.4 ± 2.5	$0. \pm .5$	$2.2 \pm .5$	$0. \pm 0.3$	$-.4 \pm .15$	$1.1 \pm .3$	$7.4 \pm .7$	$-6. \pm .7$